arXiv:1703.03242v1 [math.NT] 9 Mar 2017

On minimal additive complements of integers

Sándor Z. Kiss,[1*] Csaba Sándor,[1†] and Quan-Hui Yang[2‡]

1. Institute of Mathematics, Budapest University of Technology and Economics, H-1529 B.O. Box, Hungary

2. School of Mathematics and Statistics, Nanjing University of Information Science and Technology, Nanjing 210044, China

Abstract

Let $C, W \subseteq \mathbb{Z}$. If $C + W = \mathbb{Z}$, then the set C is called an additive complement to W in $\mathbb{Z}$. If no proper subset of C is an additive complement to W, then C is called a minimal additive complement. Let $X \subseteq \mathbb{N}$. If there exists a positive integer T such that $x+T \in X$ for all sufficiently large integers $x \in X$, then we call X eventually periodic. In this paper, we study the existence of a minimal complement to W when W is eventually periodic or not. This partially answers a problem of Nathanson.

2010 Mathematics Subject Classifications: Primary 11B13, 11B34.

Key words and phrases: Additive complements, minimal complements, periodic sets.

*Email: kisspest@cs.elte.hu. This research was supported by the National Research, Development and Innovation Office NKFIH Grant No. K115288.

†Email: csandor@math.bme.hu. This author was supported by the OTKA Grant No. K109789. This paper was supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences.

‡Email: yangquanhui01@163.com. This author was supported by the National Natural Science Foundation for Youth of China, Grant No. 11501299, the Natural Science Foundation of Jiangsu Province, Grant Nos. BK20150889, 15KJB110014 and the Startup Foundation for Introducing Talent of NUIST, Grant No. 2014r029.

1 Introduction

Let $\mathbb{N}$ denote the set of nonnegative integers and $\mathbb{Z}$ be the set of integers. For $A, B \subseteq \mathbb{Z}$ and $k \in \mathbb{Z}$, let $A + B = \{a + b : \ a \in A, \ b \in B\}$ and $kA = \{ka : a \in A\}$. If $A + B = \mathbb{Z}$, then A is called an additive complement to B in $\mathbb{Z}$. If no proper subset of A is a complement to B, then A is called a minimal complement to B in $\mathbb{Z}$.

It is easy to see that if $A \subseteq \mathbb{Z}$ is a (minimal) complement to $B \subseteq \mathbb{Z}$, then A is also a (minimal) complement to $B + d$, $d \in \mathbb{Z}$, where $B + d = \{b + d : b \in B\}$.

In 2011, Nathanson [4] proved the following theorem.

Nathanson's theorem (See [4, Theorem 8]). *Let W be a nonempty, finite set of integers. In $\mathbb{Z}$, every complement to W contains a minimal complement to W.*

In the same paper, Nathanson also posed the following problem.

Problem (See [4, Problem 11]). Let W be an infinite set of integers. Does there exist a minimal complement to W? Does there exist a complement to W that does not contain a minimal complement?

In 2012, Chen and Yang [1] answered Nathanson's problem in some aspects.

Theorem A (See [2, Theorem 1]). *Let W be a set of integers with* $\inf W = -\infty$ *and* $\sup W = +\infty$. *Then there exists a minimal complement to W.*

By Theorem A, now we only need to consider the cases $\inf W > -\infty$ or $\sup W < +\infty$. Without loss of generality, we may assume that $\inf W > -\infty$.

Theorem B (See [2, Theorem 2]). *Let $W = \{1 = w_1 < w_2 < \cdots\}$ be a set of integers and*

$$\overline{W} = (\mathbb{Z} \cap (0, +\infty)) \setminus W = \{\overline{w_1} < \overline{w_2} < \cdots\}.$$

(a) If $\limsup_{i \to +\infty}(w_{i+1} - w_i) = +\infty$, *then there exists a minimal complement to W.*

(b) If $\lim_{i\to+\infty}(\overline{w_{i+1}}-\overline{w_i})=+\infty$*, then there does not exist a minimal complement to* W*.*

Let $W=\cup_{k=0}^{\infty}[10^k,2\times10^k]$. Then it is clear that both $\limsup_{i\to+\infty}(w_{i+1}-w_i)=+\infty$ and $\limsup_{i\to+\infty}(\overline{w_{i+1}}-\overline{w_i})=+\infty$ hold. Hence $\lim_{i\to+\infty}(\overline{w_{i+1}}-\overline{w_i})=+\infty$ in Theorem B (b) cannot be changed to $\limsup_{i\to+\infty}(\overline{w_{i+1}}-\overline{w_i})=+\infty$.

In this paper, we will give further results on Nathanson's problem and deal with some sets W do not satisfy the conditions of Theorem B.

First we give some definitions. Let $S\subseteq\mathbb{N}$. Denote by $S \mod m$ the set of residues of S modulo m, i.e.,

$$S \mod m=\{r: r\in\mathbb{Z}_m, r\equiv s \pmod m \text{ for some } s\in S\}.$$

Let $X\subseteq\mathbb{N}$. If there exists a positive integer T such that $x+T\in X$ for all $x\in X$, then we call X *periodic with period* T. If $X\cup C$ is a periodic set for some finite set $C\subseteq\mathbb{N}$, then we call X *quasiperiodic*. If there exists a positive integer T such that $x+T\in X$ for all sufficiently large integers $x\in X$, then we call X *eventually periodic with period* T. Clearly, a periodic set must be quasiperiodic and a quasiperiodic set must be eventually periodic. If W is eventually periodic with $|\mathbb{N}\setminus W|=+\infty$, then both $\lim_{i\to+\infty}(w_{i+1}-w_i)<+\infty$ and $\lim_{i\to+\infty}(\overline{w_{i+1}}-\overline{w_i})<+\infty$ hold. Hence W does not satisfy the conditions of Theorem B.

For a positive integer m, we denote the set of residues mod m by $\mathbb{Z}_m$. Suppose that W is an eventually periodic set and m is the least positive period. Without loss of generality, we may assume that W has the following structure:

$$W=(m\mathbb{N}+X_m)\cup Y^{(0)}\cup Y^{(1)}, \tag{1}$$

where $X_m\subseteq\mathbb{Z}_m, (Y^{(0)} \mod m)\subseteq X_m, |Y^{(0)}|<+\infty, (Y^{(1)} \bmod m)\cap X_m=\emptyset$ and $|Y^{(1)}|<+\infty$.

In this paper, we study that what conditions are needed to ensure the

existence of a minimal complement to W. First we prove a sufficient condition.

Theorem 1. *Let W be defined in* (1). *If there exists a minimal complement to W, then there exists $C \subseteq \mathbb{Z}_m$ such that the following two conditions hold:*

(a) $C + (X_m \cup (Y^{(1)} \mod m)) = \mathbb{Z}_m$;

(b) For any $c \in C$, there exists $y \in Y^{(1)}$ such that $(c + (\{y\} \mod m)) \nsubseteq (C + X_m)$.

Remark 1. *By the proof of Theorem 1, we know that Theorem 1 also holds when $Y^{(1)}$ is an infinite set with $|Y^{(1)} \cap \mathbb{Z}^-| < +\infty$.*

Let $m = 3$, $X_m = \{0\}$, $Y^{(1)} \subseteq 3\mathbb{N} + 1$. By Theorem 1, we have the following corollary.

Corollary 1. *Let $Y \subseteq 3\mathbb{N} + 1$ and $W = 3\mathbb{N} \cup Y$. Then there does not exist a minimal complement to W.*

Remark 2. *We can choose an infinite set Y in Corollary 1 such that W is not eventually periodic. Hence, there exists an infinite, not eventually periodic set $W \subseteq \mathbb{N}$ such that $w_{i+1} - w_i \in \{1, 2, 3\}$ for all i, and there does not exist a minimal complement to W.*

Remark 3. *If $W \subseteq \mathbb{N}$ is a quasiperiodic set, then $Y^{(1)} = \emptyset$ and the condition (b) in Theorem 1 does not hold. Hence there does not exist a minimal complement to W.*

In the next step we prove a necessary condition.

Theorem 2. *Let W be defined in* (1). *Suppose that there exists $C \subseteq \mathbb{Z}_m$ such that the following two conditions hold:*

(a) $C + (X_m \cup (Y^{(1)} \mod m)) = \mathbb{Z}_m$;

(b) For any $c \in C$, there exists $y \in Y^{(1)}$ such that $(c + (\{y\} \mod m)) \nsubseteq \big((C \setminus \{c\}) + (X_m \cup (Y^{(1)} \mod m))\big)$.

Then there exists a minimal complement to W.

By Theorems 1 and 2, we have the following corollary.

Corollary 2. *Let* $W = (m\mathbb{N} + X_m) \cup Y^{(0)} \cup \{a\}$, *where* $X_m \subseteq \mathbb{Z}_m$, $(Y^{(0)} \mod m) \subseteq X_m$ *and* $(\{a\} \mod m) \nsubseteq X_m$. *Then there exists a minimal complement to* W *if and only if there exists a subset* $C \subseteq \mathbb{Z}_m$ *such that:*

(1) $C + (X_m \cup (\{a\} \mod m)) = \mathbb{Z}_m$;

(2) For any $c \in C$, $(c + (\{a\} \mod m)) \nsubseteq (C + X_m)$.

We see that Theorems 1 and 2 transfer Nathanson's problem into a finite modulo version. In the next theorem, we give a sufficient and necessary condition, but we cannot bound the module.

Theorem 3. *Let* W *be defined in (1). There exists a minimal complement to* W *if and only if there exists* $T \in \mathbb{Z}^+, m \mid T$, *and* $C \subseteq \mathbb{Z}_T$ *such that*

(1) $C + (X_T \cup (Y^{(1)} \mod T)) = \mathbb{Z}_T$, *where* $X_T = \cup_{i=0}^{\frac{T}{m}-1}\{im + X_m\}$;

(2) for any $c \in C$, *there exists* $y \in Y^{(1)}$ *for which* $(c + (\{y\} \mod T)) \nsubseteq ((C \setminus \{c\}) + (X_T \cup (Y^{(1)} \mod T)))$.

Finally, as a complement to Remark 2, we give the following theorem.

Theorem 4. *There exists an infinite, not eventually periodic set* $W \subseteq \mathbb{N}$ *such that* $w_{i+1} - w_i \in \{1, 2\}$ *for all* i *and there exists a minimal complement to* W.

Now we pose two problems for further research.

Problem 1. *We know that Theorem 1 also holds when* $Y^{(1)}$ *is infinite. Is Theorem 2 also true when* $Y^{(1)}$ *is infinite?*

Problem 2. *Does there exist an upper bound for* T *in Theorem 3 using* $m, Y^{(0)}$ *and* $Y^{(1)}$*?*

2 Proofs

Proof of Theorem 1. Suppose that D is a minimal complement to W. For $i \in \{0, 1, \ldots, m-1\}$, let $D_i = \{d \in D : \ d \equiv i \pmod m\}$ and

$$C = \{j : \ 0 \le j \le m-1 \text{ and } |D_j \cap \mathbb{Z}^-| = +\infty\}.$$

For any $t \in \{0, 1, \ldots, m-1\} \setminus C$, the set $\{d \in D : \ d \equiv t \pmod m\} + W$ does not contain any sufficiently small negative integers. It follows from $D + W = \mathbb{Z}$ that $C + ((W \cup Y^{(1)}) \bmod m) = \mathbb{Z}_m$. That is, $C + (X_m \cup (Y^{(1)} \bmod m)) = \mathbb{Z}_m$.

Next we shall prove (2). Suppose that there exists $c \in C$ such that $(c + (\{y\} \bmod m)) \subseteq (C + X_m)$ for any $y \in Y^{(1)}$. We take an integer $d \in D$ such that $d \equiv c \pmod m$ and we shall prove that $D \setminus \{d\}$ is also a complement to W. For any integer n, write $n = d' + w$, where $d' \in D$ and $w \in W$.

Case 1. Assume $d' \neq d$. Then $n = d' + w \in (D \setminus \{d\}) + W$.

Case 2. Assume $d' = d$.

Subcase 2.1. $\{w \bmod m\} \subseteq X_m$. In this case, there exists a positive integer k_0 such that $w + km \in W$ for all integers $k \geq k_0$. Since $|D_c \cap \mathbb{Z}^-| = +\infty$, it follows that there exists an integer $k \geq k_0$ such that $d - km \in D$. Hence $n = (d - km) + (w + km)$, where $d - km \in D \setminus \{d\}$ and $w + km \in W$. That is, $n \in (D \setminus \{d\}) + W$.

Subcase 2.2. $w \in Y^{(1)}$. Since $(c + (\{y\} \bmod m)) \subseteq C + X_m$ for any $y \in Y^{(1)}$ and $d \equiv c \pmod m$, $w \in Y^{(1)}$, it follows that $\{n \bmod m\} = \{d + w \bmod m\} \subseteq C + X_m$. Hence there exist a $c' \in D$ with $c' \pmod m \in C$ and $x \in W$ with $x \bmod m \in X_m$ such that $n \equiv \ c' + x \pmod m$. We choose a sufficiently large integer k such that $c' - km \in D$, $c' - km \neq d$ and $x + km \in W$. Hence $n = (c' - km) + (x + km)$, where $c' - km \in D \setminus \{d\}$ and $x + km \in W$.

Hence, $(D \setminus \{d\}) + W = \mathbb{Z}$ which contradicts the fact that D is a minimal complement. Therefore, (2) holds. □

Proof of Theorem 2. Let $C_1 = C + X_m$, $C_2 = \mathbb{Z}_m \setminus C_1$,

$$C' = \{d \in \mathbb{Z} : \ d \equiv c \pmod m \text{ for some } c \in C\},$$

$$C_1' = \{d \in \mathbb{Z} : \ d \equiv c \pmod m \text{ for some } c \in C_1\},$$

$$C_2' = \{d \in \mathbb{Z} : \ d \equiv c \pmod m \text{ for some } c \in C_2\}.$$

By (1), we have $C' + W = \mathbb{Z}$. Since $C + X_m = C_1$, it follows that

$$\big(C' + (W \setminus Y^{(1)})\big) \bmod m = C + X_m = C_1,$$

and so $\big(C' + (W \setminus Y^{(1)})\big) \cap C_2' = \emptyset$. Noting that $C' + W = \mathbb{Z}$, we have $C' + Y^{(1)} \supseteq C_2'$. Since $Y^{(1)}$ is a finite set, by the proof of Nathanson's theorem (See [4, Theorem 4, page 2015]), there exists $D' \subseteq C'$ such that $D' + Y^{(1)} \supseteq C_2'$ and for any $d \in D'$,

$$(D' \setminus \{d\}) + Y^{(1)} \not\supseteq C_2'.$$

Next we shall prove that D' is a minimal complement to W.

For $i \in C$, let $D_i' = \{d \in D' : d \equiv i \pmod m\}$. First we prove that $|D_i' \cap \mathbb{Z}^-| = +\infty$ for all $i \in C$. Suppose that there exists a $j \in C$ such that $|D_j' \cap \mathbb{Z}^-| < +\infty$. By (2), there exists a $y \in Y^{(1)}$ such that

$$\big(j + (\{y\} \bmod m)\big) \not\subseteq (C \setminus \{j\}) + (X_m \cup Y^{(1)}),$$

and so

$$D' + Y^{(1)} \not\supseteq \{d \in \mathbb{Z} : \{d \bmod m\} = j + \{y\} \bmod m\}.$$

Noting that $\big(j+(\{y\} \bmod m)\big) \not\subseteq C+X_m = C_1$, we have $\big(j+(\{y\} \bmod m)\big) \subseteq C_2$. It follows that $D'+Y^{(1)} \not\supseteq C_2'$, a contradiction. Hence, $|D_i' \cap \mathbb{Z}^-| = +\infty$ for all $i \in C$.

Next we prove that $D^{'}$ is a complement. For any integer $n \in C_1'$, by $C+X_m = C_1$, there exists $c \in C$ and $x \in X_m$ such that $\{n\} \bmod m = c+x$. Since $|D_c' \cap \mathbb{Z}^-| = +\infty$, there exists a sufficiently small negative integer $d \in D_c'$ such that $n - d > 0$, and so

$$n = d + (n - d) \in D_c' + (m\mathbb{N} + X_m) \subseteq D' + W.$$

Hence $C_1' \subseteq D' + W$. On the other hand, $D' + W \supseteq D' + Y^{(1)} \supseteq C_2'$. Therefore, $D' + W = \mathbb{Z}$.

Finally, we prove that $D^{'}$ is a minimal complement. For any $d \in D'$, by $(D' \setminus \{d\}) + Y^{(1)} \not\supseteq C_2'$ and

$$\big((D' \setminus \{d\}) + (W \setminus Y^{(1)})\big) \bmod m = C + X_m = C_1,$$

we have

$$((D' \setminus \{d\}) + (W \setminus Y^{(1)})) \cap C_2' = \emptyset,$$

and so $(D' \setminus \{d\}) + W \not\supseteq C_2'$. Hence $(D' \setminus \{d\}) + W \neq \mathbb{Z}$.

Therefore, D' is a minimal complement to W. □

Proof of Theorem 3. Assume that the set W satisfies the conditions of Theorem 3. Applying Theorem 2 with $m = T$, it follows that W has a minimal complement.

Suppose that W has a minimal complement denoted by E. We will prove the existence of a positive integer T and a set $C \subseteq \mathbb{Z}_T$ which satisfy the conditions of Theorem 3. We will show that there exist positive integers K and L with $L > K$ such that $T = L - K$. We will prove that this integer T and the set

$$C = \{l \bmod (L-K) : K \leq l < L,\ l \in E\}$$

are suitable.

For $0 \leq i < m$, let

$$E_i^- = \{e : e < 0, e \in E, e \equiv i \bmod m\}.$$

Let $0 \leq i_1 < i_2 < \ldots < i_t < m$ be the sequence of indices with $|E_{i_j}^-| = \infty$. It is clear that there exists an integer N_0 such that, $e \in E$ and $e \leq N_0$ imply that $e \in E_{i_j}$ for some i_j. Let

$$y_+ = \max\{y : y \in Y^{(0)} \cup Y^{(1)}\},$$

$$y_- = \min\{y : y \in Y^{(0)} \cup Y^{(1)}\},$$

and $y_0 = y_+ - y_- + 1$. Let $\chi_E(k)$ denote the characteristic function of the set E, i.e.,

$$\chi_E(k) = \begin{cases} 1, \text{ if } k \in E; \\ 0, \text{ if } k \notin E. \end{cases}$$

Put $y^{(-)} = -y_+ + \min\{0, y_-\}$, $y^{(+)} = -y_- + \max\{0, y_+\}$ and define the positive integer A by $A = N_0 + \min\{0, y_-\}$. Consider the following vectors:

$$\begin{aligned}
\mathbf{v}_A &= (\chi_E(A + y_-), \chi_E(A + y_- + 1), \ldots, \chi_E(A + y_+)) , \\
\mathbf{v}_{A-m} &= (\chi_E(A - m + y_-), \chi_E(A - m + y_- + 1), \ldots, \chi_E(A - m + y_+)) , \\
&\vdots \\
\mathbf{v}_{A-im} &= (\chi_E(A - im + y_-), \chi_E(A - im + y_- + 1), \ldots, \chi_E(A - im + y_+)) , \\
&\vdots
\end{aligned}$$

It is clear that there are infinitely many vectors $\mathbf{v}_A, \mathbf{v}_{A-m}, \ldots, \mathbf{v}_{A-im}, \ldots$, each of them has y_0 coordinates, which are 0 or 1. Since there are at most 2^{y_0} different vectors, by the pigeon hole principle, there exists a vector $\mathbf{v}$ among them which occurs infinitely many times. In other words there exists an infinite sequence $0 \leq k_1 < k_2 < \cdots$ such that $\mathbf{v}_{A-k_i m} = \mathbf{v}$. Define L by $L = A - k_1 m$. Obviously it can be chosen a k_i large enough such that

$$[A - k_i m, A - k_1 m[\cap E_{i_j} \neq \emptyset$$

holds for every index i_j and $k_i m - k_1 m \geq y_0$. In view of this fact we define K by $K = A - k_i m$ and $T = L - K$. It follows from the definition that K and L have the following properties.

$$L \leq N_0 + \min\{0, y_-\}, \tag{2}$$

$$K \leq L - y_0, \tag{3}$$

$$m \mid L - K, \tag{4}$$

$$\chi_E(K + i) = \chi_E(L + i), \quad for \ \ y^{(-)} \leq i \leq y^{(+)}, \tag{5}$$

$$[K, L[\cap E_{i_j} \neq \emptyset \text{ for all } i_j. \tag{6}$$

In the next step we show that the positive integer T and set C defined above satisfy the conditions of Theorem 3.

We know from the conditions of Theorem 3 that $W = (T\mathbb{N}+X_T)\cup Y^{(0)}\cup Y^{(1)}$ and $X_T \subseteq \mathbb{Z}_T$. First we prove that $C + (X_T \cup (Y^{(1)} \bmod T)) = \mathbb{Z}_T$. Let $K \leq l < L$. It follows that $l = e + w$, where $e \in E$ and $w \in W$. As $w \geq \min\{0, y_-\}$, it follows from (1) that $e = l - w < L - \min\{0, y_-\} \leq N_0$, thus we have $e \in E_{i_j}^-$.

Suppose that $w \in Y^{(0)}\cup Y^{(1)}$. Then we have $y_- \leq w \leq y_+$, which implies that $e = l - w$, where $K - y_+ \leq e < L - y_-$. We have three cases.

Case 1. $K - y_+ \leq e < K$. By (5) we have $e + (L - K) \in E$ and $L - y_+ \leq e + L - K < L$. Thus we have $l \equiv c + w \pmod T$, where $c \in C$ and $w \in X_T \cup (Y^{(1)} \bmod T)$.

Case 2. $K \leq e < L$. It follows that $l \equiv c+w \pmod T$, where $c \in C$ and $w \in X_T \cup (Y^{(1)} \bmod T)$.

Case 3. $L \leq e < L - y_-$. By (5) we have $e - (L - K) \in E$ and $K \leq e - (L - K) < K - y_-$, thus we have $l \equiv c + w \pmod T$, where $c \in C$ and $w \in X_T \cup (Y^{(1)} \bmod T)$.

Suppose that $w \in T\mathbb{N} + X_T$. Since $w \geq 0$ and $e = l - w < L \leq N_0$, we have $e \in E_{i_j}^-$, which implies that $e \equiv i_j \pmod m$. It follows from (6) that there exists an $e^{'}$ such that $e^{'} \in E_{i_j}^-$ and $K \leq e^{'} < L$, thus we have $e^{'} \equiv i_j \pmod m$. Let $w \equiv x \pmod m$, where $0 \leq x < m$, $x \in X_m$. Obviously, $l \equiv e+w \equiv i_j+x \equiv e^{'}+x \pmod m$. By using (4) it follows that there exists a u with $0 \leq u < \frac{L-K}{m}$ such that $l \equiv e^{'} + um + x \pmod{L - K}$. Therefore, $c \equiv e^{'} \pmod{L-K}$, where $0 \leq c < L-K$, $c \in C$ and $0 \leq um+x < L-K$ and $um + x \in X_T$ are suitable.

In the next step we show that the second condition of Theorem 3 holds. For every $K \leq e < L \leq N_0$ and $e \in E$, there exists a $w \in W$ such that $e + w \neq e^{'} + w^{'}$, when $e \neq e^{'}$. If $w \in (m\mathbb{N} + X_m) \cup Y^{(0)}$, then there exists a positive integer s such that $e + w = (e - sm) + (w + sm)$, where $e - sm \in E$ and $w + sm \in W$, which is absurd. Then we may assume

that $w \in Y^{(1)}$. It follows that $K + y_- \leq e + w \leq L + y_+$. Obviously, it is enough to prove that $e + w \not\equiv e^{'} + w^{'} \pmod{L-K}$, where $K \leq e^{'} < L$, $e^{'} \in E$, $e \neq e^{'}$ and $w^{'} \in X_T \cup (Y^{(1)} \bmod T)$. Suppose that for $w^{'} \in X_T$ we have $e + w = e^{'} + w^{'} + t(L-K)$ for some integer t. Then we have $e + w = e^{'} + w^{'} + t(L-K) = (e^{'} - sm) + (w^{'} + sm + t(L-K))$, where $e^{'} - sm \in E$ and $w^{'} + sm + t(L-K) \in W$ for some positive integer s, which is absurd.

For any $w^{'} \in Y^{(1)}$, clearly we have $K + y_- \leq e + w, e^{'} + w^{'} \leq L + y_+$. Assume that $e+w \equiv e^{'}+w^{'} \pmod{L-K}$. It follows that either $e+w = e^{'}+w^{'}$ or $e + w = e^{'} + w^{'} + (L-K)$ or $e + w = e^{'} + w^{'} - (L-K)$.

Case 1. $e+w = e^{'}+w^{'}-(L-K)$. Then we have $L+y_- \leq e^{'}+w^{'} \leq L+y_+$. Thus we have $L + y_- - y_+ \leq e^{'} \leq L + y_+ - y_-$. It follows from (5) that $e^{'} - (L-K) \in E$ which implies that $e + w = (e^{'} - (L-K)) + w^{'}$. Therefore we have $w = w^{'}$ and $e = e^{'} - (L-K)$, which is a contradiction because $K \leq e, e^{'} < L$.

Case 2. $e + w = e^{'} + w^{'}$. Then we have $e = e^{'}$ and $w = w^{'}$, which is impossible.

Case 3. $e+w = e^{'}+w^{'}+(L-K)$. Then we have $K+y_- \leq e^{'}+w^{'} \leq K+y_+$. Thus we have $K + y_- - y_+ \leq e^{'} \leq K + y_+ - y_-$. It follows from (5) that $e^{'} + (L-K) \in E$ which implies that $e + w = (e^{'} + (L-K)) + w^{'}$. Therefore we have $w = w^{'}$ and $e = e^{'} + (L-K)$ which is absurd because $K \leq e, e^{'} < L$.

The proof of Theorem 3 is completed. □

Proof of Theorem 4. By induction we can construct $\{d_i\}_{i=1}^{\infty}$, $\{W_i\}_{i=1}^{\infty}$ and $\{c_i\}_{i=1}^{\infty}$ such that

(i) $d_1 = -1,\ W_1 = \{1, 2, \ldots, 12\},\ c_1 = -3$;

(ii) d_i is the largest negative integer $\notin W_{i-1} + \{c_1, c_2, \ldots, c_{i-1}\}$ for $i \geq 2$;

(iii) take an integer $c_i < d_i + 2c_{i-1}$ for all $i \geq 2$;

(iv) for $i \geq 2$, let $W_i = W_{i-1} \cup \left([-2c_{i-1}, -2c_i + 1] \setminus \cup_{j=1}^{i-1}\{-c_i + d_j\}\right)$.

Let $W = \cup_{i=1}^{\infty} W_i$ and $C = \{c_i\}_{i=1}^{\infty}$.

Now we prove that C is a minimal complement to W.

First we prove $d_{i+1} - d_i \leq -2$ for all integers $i \geq 1$. Clearly $d_2 = -3$, $d_2 - d_1 = -2$. Suppose that $d_{i+1} - d_i \leq -2$ for all integers $i < k$ ($k \geq 2$). Since

$$d_k = (d_k - c_k) + c_k, \quad d_k - 1 = (d_k - 1 - c_k) + c_k,$$

$$-2c_{k-1} \leq d_k - 1 - c_k < d_k - c_k < -c_k + d_{k-1},$$

it follows that $d_k - c_k,\ d_k - 1 - c_k \in W_k$ and then $d_k, d_k - 1 \in W_k + \{c_k\}$. Hence $d_{k+1} \leq d_k - 2$. By (iv), we have $w_{k+1} - w_k \in \{1, 2\}$. Since $d_k \to -\infty$, by (ii) we have $(-\infty, 9] \subseteq W + C$. For any integer $n \geq 10$, there exists an i such that $-c_{i-1} \leq n < -c_i$. Hence

$$-c_i + d_1 < -c_{i-1} - c_i \leq n - c_i < -2c_i,$$

and so $n - c_i \in W_i$, that is, $n \in W_i + \{c_i\}$. Therefore, $W + C = \mathbb{Z}$.

Next, we prove that the complement C is minimal. For any positive integer i, we write $d_i = c + w$ with $c \in C$ and $w \in W$. Now we shall prove that $c = c_i$. By (iv), we have $d_i - c_j \notin W$ for all integers $j > i$. Hence $c \neq c_j$ for all integers $j > i$. Since $-2c_{i-1}$ is the minimal value of $W \setminus W_{i-1}$ and for any positive integers $j \leq i - 1$, $d_i - c_j \leq d_i - c_{i-1} < -2c_{i-1}$, it follows that $d_i - c_j \notin W \setminus W_{i-1}$ for all positive integer $j \leq i - 1$. Noting that $d_i \notin W_{i-1} + \{c_1, \ldots, c_{i-1}\}$, we have $d_i \notin W + \{c_1, c_2, \ldots, c_{i-1}\}$. Hence $c = c_i$.

Therefore, C is a minimal complement to W. Furthermore, by (iii), we can choose suitable c_i such that W is infinite and not eventually periodic. □

3 Acknowledgement

This work was done during the third author visiting to Budapest University of Technology and Economics. He would like to thank Dr. Sándor Kiss and Dr. Csaba Sándor for their warm hospitality.

References

[1] Y.-G. Chen, J.-H. Fang, *On additive complements, II,* Proc. Amer. Math. Soc. 139 (3) (2011) 881-883.

[2] Y.-G. Chen, Q.-H. Yang, *On a problem of Nathanson related to minimal additive complements* SIAM J. Discrete Math. 26 (4) (2012) 1532-1536.

[3] J.-H. Fang, Y.-G. Chen, *On additive complements,* Proc. Amer. Math. Soc. 138 (6) (2010) 1923-1927.

[4] M.-B. Nathanson, *Problems in additive number theory, IV: Nets in groups and shortest length g-adic representations,* Int. J. Number Theory 7 (3) (2011) 1999-2017.

[5] I. Z. Ruzsa, *On the additive completion of linear recurrence sequences,* Periodica Math. Hungar 9 (1978) 285-291.

[6] I. Z. Ruzsa, *Additive completion of lacunary sequences,* Combinatorica 21 (2) (2001) 279-291.